



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Even Abundant Numbers.

BY L. E. DICKSON.

1. THEOREM. *There is only a finite number of primitive non-deficient numbers having a given number n of distinct odd prime factors and a given number m of factors 2.*

Let p_1, \dots, p_n be primes in ascending order. Then

$$a = 2^m p_1^{e_1} p_2^{e_2} \dots p_n^{e_n}$$

is deficient if

$$G \cdot \left(\frac{p_1}{p_1 - 1} \right)^n < 2, \quad G \equiv \frac{2^{m+1} - 1}{2^m},$$

and hence if $p_1 > r/(r-1)$, where r is the positive real n -th root of $2/G$, whence $r > 1$. Thus the least odd prime factor p_1 of a non-deficient number is limited. To prove by induction a like result for each p_i , let p_1, \dots, p_ν be given primes, and $\nu < n$. The divisor

$$\alpha = 2^m p_1^{e_1} \dots p_\nu^{e_\nu}$$

of a primitive a must be deficient. By Lemma B of the former paper, each deficient α is a product of some one of a finite number of α 's by a number having no prime factor other than p_1, \dots, p_ν . We may now complete the proof as in the former paper, inserting in Π_μ the additional factor G .

For example, the primitive non-deficient $2^m p^e$ are $2^m p$, where $2^m - 1 < p \leq 2^{m+1} - 1$.

2. In the determination of all even non-deficient numbers not exceeding a given number L , use may be made of the following theorems.

THEOREM. *Let $2^l \leq L < 2^{l+1}$ and $\lambda = l/2$ or $(l+1)/2$, according as l is even or odd. If n is any integer for which $l > n \geq \lambda$ and k is any odd integer for which $2^n k \leq L$, then $2^n k$ is non-deficient.*

We have $2^\lambda k \leq 2^n k \leq L < 2^{l+1}$. Now $l+1 = 2\lambda+1$ or 2λ , according as l is even or odd. Thus $k < 2^{n+1}$ or 2^n , in the respective cases. In each case, $k \leq N$, where $N = 2^{n+1} - 1$. Thus k has a prime factor $p \leq N$. Since $2^n p$ is non-deficient, the same is true of its multiple $2^n k$.

3. THEOREM. *Let ν be the least integer for which*

$$2^\nu (2^{\nu+1} - 1)^2 \geq L,$$

and n any integer $\geq \nu$. Let c be any odd composite number for which $2^n c < L$. Then $2^n c$ is abundant.

Set $N = 2^{n+1} - 1$. Then $2^n N^2 \geq L$ and $c < N^2$. Hence the composite number c has a prime factor $p < N$. Since $2^n p$ is abundant, its multiple $2^n c$ is abundant.

This theorem, in connection with the fact that $2^n p$ (p an odd prime) is abundant if and only if $p < N$, enables one to write down immediately all abundant numbers $2^n k < L$, where $n \geq \nu$ (cf. § 8).

4. THEOREM. Set $s = 2^{r+1} - 1$, $t = 2^r - 1$. The non-deficient even numbers $2^{r-1}k$ less than* $2^r s^2$ are those in which k has a prime factor $\leq t$ and those in which k is a product of two primes $t + P$ and $t + Q$, where $0 < P < t$, $P < Q \leq (t^2 + t)/P$. If $P = t - 1$, then $Q = t + 1$.

Let every prime factor of k exceed t . Since

$$(2^r + 1)^3 \geq 2s^2$$

for every $r > 0$, k can not have three prime factors. If p is an odd prime, $2^n p^e$ is deficient if

$$\frac{2^{n+1} - 1}{2^n} \frac{p}{p-1} < 2, \quad p > 2^{n+1}$$

Hence $2^{r-1}p^e$ is deficient if $p > t$. Finally, if p, q are distinct primes $> t$ and $q > p$, then $2^{r-1}pq$ is non-deficient if and only if

$$pq \leq t(p + q + 1).$$

Set $p = t + P$, $q = t + Q$. Then the condition becomes $PQ \leq t^2 + t$. But $Q \geq P + 2$. Hence $P < t$. If $P = t - 1$, then $Q < t + 3$, so that $Q = t + 1$. For, if $Q \geq t + 3$,

$$PQ \geq t^2 + t + t - 3, \quad PQ > t^2 + t.$$

Indeed, $r \geq 2$, $t \geq 3$; while if $t = 3$ and if $Q = t + 3$, then $q = 9$.

5. THEOREM. The non-deficient numbers $2^{r-2}k$ less than $2^r s^2$ and having† $r > 3$ are those in which k has a prime factor $\leq r$, where $r = 2^{r-1} - 1$, those in which k is the product of two primes $r + P$ and $r + Q$, where $0 < P < r$, $P < Q \leq (r^2 + r)/P$, and the following abundant numbers:

$$\begin{aligned} &2^4 37 \cdot 41^2, \quad 2^4 37^2 q (q = 41, 43, 47), \quad 2^3 17 \cdot 19 l (23 \leq l \leq 47), \\ &2^3 17 \cdot 23 l (l = 29, 31, 37), \quad 2^3 17 \cdot 29 \cdot 31, \quad 2^3 19 \cdot 23 l (l = 29, 31), \\ &2^3 17^2 q (q < 55), \quad 2^3 19^2 q (q < 45), \quad 2^3 23^2 q (q < 31), \quad 2^3 29^2 17, \\ &2^2 11^2 q (q < 26), \quad 2^2 13^2 q (q < 19), \quad 2^2 17^2 q (q < 14), \\ &2^2 11 \cdot 13 l (l = 17, 19, 23), \quad 2^2 11 \cdot 17 \cdot 19, \end{aligned}$$

where q and l are primes, $q > 7$ in the fourth line, $q > 15$ in the third line.

Let every prime factor of k exceed r . Since

$$(2^{r-1} + 1)^4 > 2^{4r-4} \geq 4 \cdot 2^{2r+2} > 4s^2$$

* Including all $< L$, since $L \leq 2^r s^2$ by § 3.

† Those with $r = 3$ are given in § 7.

for $\nu > 3$, k can not have four prime factors. Since

$$(2^{\nu-1} + 1)^3 > 2^{3\nu-3} \geq 4 \cdot 2^{2\nu+2} > 4s^2$$

for $\nu \geq 7$, k has at most two prime factors unless $\nu = 6, 5$ or 4 . As in § 4, $2^{\nu-2}p^e$ is deficient if $p > r$, since then $p > 2^{\nu-1}$. The case in which k is the product of two distinct prime factors may be treated as in § 4. We shall next consider $2^{\nu-2}p^2q$, where $\nu = 6, 5$ or 4 , and p, q are distinct primes $> r$. First, let $\nu = 6$, whence $r = 31$, $s = 127$. The least p is 37; by $37^2q < 4s^2$, $q < 48$; 2^437^2q is abundant if $q < 229$. For $p = 41$, $4s^2/p^2 < 39$, whence $q = 37$. Next, let $\nu = 5$, whence $r = 15$, $s = 63$. For $p = 17, 19, 23, 29, 31$, $p^2q < 4s^2$ for $q < 55, 45, 31, 19, 17$, respectively, the final p being therefore excluded; while 2^3p^2q is abundant for $q < 243, 94, 50, 34, 31$, respectively. Finally, let $\nu = 4$, whence $r = 7$, $s = 31$. For $p = 11, 13, 17, 19$, $p^2q < 4s^2$ for $q < 32, 23, 14, 11$; while 2^2p^2q is abundant for $q < 26, 19, 14, 13$.

It remains only to consider $2^{\nu-2}pql$, where p, q, l are distinct primes $> r$, arranged in ascending order. First, let $\nu = 6$. If $p \geq 41$, $pql > 41^3 > 4s^2$. Hence $p = 37$. But $37 \cdot 41 \cdot 43 > 4s^2$. For $\nu = 5$ or 4 , the numbers $< 4s^2$ are listed in the theorem, all being abundant.

6. While we might treat similarly the cases $\nu = 3$, etc., the results already obtained, together with those in § 7 for non-deficient numbers $2k$, enable us to tabulate in § 8 the even non-deficient numbers less than $2^\nu s^2$ for $\nu = 4$, namely, $< 2^431^2 = 15376$. Indeed, under this limit L , there is no primitive non-deficient number $2k$, where k is an odd number with more than three distinct prime factors (a case not treated in § 7). First, 3 is not a factor. If 5 is a factor, 7 is not, and $2k \geq 2 \cdot 5 \cdot 11 \cdot 13 \cdot 17 = 24310 > L$. If 5 is not a factor, $2k \geq 2 \cdot 7 \cdot 11 \cdot 13 \cdot 17$, which exceeds the preceding.

7. THEOREM. *The primitive non-deficient numbers $2k$, where k is an odd number with at most three distinct prime factors, are:*

$$\begin{aligned} &2 \cdot 3, 2 \cdot 5 \cdot 7, 2 \cdot 5^2 \cdot 11, 2 \cdot 5^2 \cdot 13, 2 \cdot 5 \cdot 11 \cdot p (13 \leq p \leq 53), \\ &2 \cdot 5 \cdot 11^2 \cdot p (59 \leq p \leq 89, 2 \cdot 5 \cdot 11^3 \cdot 97, 2 \cdot 5 \cdot 13 \cdot p (17 \leq p \leq 31), \\ &2 \cdot 5 \cdot 13^2 \cdot 37, 2 \cdot 5 \cdot 17 \cdot 19, 2 \cdot 5^2 \cdot 17 \cdot p (23 \leq p \leq 61), 2 \cdot 5^2 \cdot 17^2 \cdot p (67 \leq p \leq 79), \\ &2 \cdot 5^2 \cdot 17^3 \cdot 83, 2 \cdot 5^3 \cdot 17 \cdot p (67 \leq p \leq 109), 2 \cdot 5^3 \cdot 17^2 \cdot p (113 \leq p \leq 173), \\ &2 \cdot 5^3 \cdot 17^3 \cdot 179, 2 \cdot 5^3 \cdot 17^3 \cdot 181^2, 2 \cdot 5^3 \cdot 17^4 \cdot 181, 2 \cdot 5^4 \cdot 17 \cdot p (p = 113, 127), \\ &2 \cdot 5^4 \cdot 17^2 \cdot p (179 \leq p \leq 223), 2 \cdot 5^4 \cdot 17^3 \cdot p (p = 227, 229, 233), 2 \cdot 5^5 \cdot 17 \cdot 131, \\ &2 \cdot 5^5 \cdot 17^2 \cdot p (p = 227, 229, 233), 2 \cdot 5^5 \cdot 17^2 \cdot 239^2, 2 \cdot 5^5 \cdot 17^3 \cdot p (p = 239, 241), \\ &2 \cdot 5^5 \cdot 17^3 \cdot 251^2, 2 \cdot 5^6 \cdot 17^2 \cdot p (p = 239, 241), 2 \cdot 5^6 \cdot 17^3 \cdot 251, \\ &2 \cdot 5^2 \cdot 19 \cdot p (23 \leq p \leq 43), 2 \cdot 5^2 \cdot 19 \cdot 47^2, 2 \cdot 5^2 \cdot 19^2 \cdot p (p = 47, 53), \\ &2 \cdot 5^3 \cdot 19 \cdot p (47 \leq p \leq 61), 2 \cdot 5^3 \cdot 19 \cdot 67^2, 2 \cdot 5^3 \cdot 19^2 \cdot p (67 \leq p \leq 79), \end{aligned}$$

$2 \cdot 5^3 19^3 83^2$, $2 \cdot 5^4 19 p$ ($p = 67, 71, 73$), $2 \cdot 5^4 19^2 p$ ($p = 83, 89$),
 $2 \cdot 5^2 23 p$ ($p = 29, 31$), $2 \cdot 5^3 23 p$ ($p = 37, 41$), $2 \cdot 5^3 23^2 43$, $2 \cdot 5^4 23^2 47$,
 $2 \cdot 5^3 29 \cdot 31^2$, $2 \cdot 5^3 29^2 31$, $2 \cdot 5^4 29 \cdot 31$,
 $2 \cdot 7 \cdot 11 \cdot 13$, $2 \cdot 7^2 13^2 17$, $2 \cdot 7^3 13 \cdot 17^2$, $2 \cdot 7^3 13^3 19^3$, $2 \cdot 7^3 13^4 19^2$, $2 \cdot 7^4 13^2 19^3$,
 $2 \cdot 7^4 13^3 19^2$.

The proof is similar to that used in the former paper.

8. We are now in a position to give the even non-deficient numbers* $< L = 15000$. This L is just under the maximum limit given by $\nu = 4$ (§ 6). Moreover, this L was the convenient limit used in listing the primitive odd abundant numbers in the former paper.

The non-deficient numbers $2^n k < L$, with $n \geq 4$, are (§§ 2, 3):

$2^{12} 3$, $2^{11} k$ ($k \leq 7$), $2^{10} k$ ($k \leq 13$), $2^9 k$ ($k \leq 29$), $2^8 k$ ($k \leq 57$), $2^7 k$ ($k \leq 117$),
 $2^6 c$ ($c \leq 233$), $2^6 p$ ($p \leq 127$), $2^5 c$ ($c \leq 467$), $2^5 p$ ($p < 63$), $2^4 c$ ($c \leq 937$), $2^4 p$ ($p \leq 31$),
 where k, c, p are odd and > 1 , c is composite and p prime.

The non-deficient $2^3 k < L$ are (§ 4) those with k having a prime factor ≤ 13 ($k \leq 1875$) and $k = 29 \cdot 31$, $23 q$ ($29 \leq q \leq 43$), $19 q$ ($23 \leq q \leq 73$), $17 q$ ($19 \leq q \leq 109$). In the last case only the limit for abundance ($q \leq 131$) exceeded the limit required by L .

The non-deficient $2^2 k < L$ are (§ 5) those with $k < 3750$ having a factor 3, 5 or 7; those with $k = 11 q$ ($q = 13, 17, 19$); and

$2^2 11^2 q$ ($q = 13, 17, 19, 23$), $2^2 13^2 q$ ($q = 11, 17$), $2^2 17^2 q$ ($q = 11, 13$),
 $2^2 11 \cdot 13 l$ ($l = 17, 19, 23$), $2^2 11 \cdot 17 \cdot 19$.

The primitive non-deficient $2 k < L$ are (§ 7) those in the first two lines of the theorem below.

The primitives are now found at once. For instance, $2^2 k$ with k having a factor 3, 5 or 7 is a multiple of one of the primitives $2 \cdot 3$, $2^2 5$, $2^2 7$. In the list of non-deficients $2^n k$, $n \geq 4$, k or c is always less than the square of the maximum prime p giving a non-deficient $2^n p$. Hence we obtain the

THEOREM. *The 98 primitive even non-deficient numbers < 15000 are:*

$2 \cdot 3$, $2 \cdot 5 \cdot 7$, $2 \cdot 5^2 11$, $2 \cdot 5^2 13$, $2 \cdot 5 \cdot 11 p$ ($13 \leq p \leq 53$),
 $2 \cdot 5 \cdot 13 p$ ($17 \leq p \leq 31$), $2 \cdot 5 \cdot 17 \cdot 19$, $2 \cdot 7 \cdot 11 \cdot 13$,
 $2^2 5$, $2^2 7$, $2^2 11 q$ ($q = 13, 17, 19$), $2^2 11^2 23$, $2^2 13 \cdot 17^2$, $2^2 13^2 17$,
 $2^3 11$, $2^3 13$, $2^3 17 q$ ($19 \leq q \leq 109$), $2^3 19 q$ ($23 \leq q \leq 73$), $2^3 23 q$ ($29 \leq q \leq 43$), $2^3 29 \cdot 31$,
 $2^4 p$ ($17 \leq p \leq 31$), $2^5 p$ ($37 \leq p \leq 61$), $2^6 p$ ($67 \leq p \leq 127$).

* All < 6232 are tabulated in *Quar. Jour. Math.*, 1913, pp. 274-7.